

Gamma Rays from Bino-like Dark Matter in the MSSM

Jason Kumar¹ and Pearl Sandick²

¹*Department of Physics and Astronomy,
University of Hawai'i, Honolulu, HI 96822, USA*

²*Department of Physics and Astronomy,
University of Utah, Salt Lake City, UT 84112, USA*

Abstract

We consider regions of the parameter space of the Minimal Supersymmetric Standard Model (MSSM) with bino-like neutralino dark matter in which a large fraction of the total dark matter annihilation cross section in the present era arises from annihilation to final states with monoenergetic photons. The region of interest is characterized by light sleptons and heavy squarks. We find that it is possible for the branching fraction to final states with monoenergetic photons to be comparable to that for continuum photons, but in those cases the total cross section will be so small that it is unlikely to be observable. For models where dark matter annihilation may be observable in the present era, the branching fraction to final states with monoenergetic photons is $\mathcal{O}(1 - 10\%)$.

PACS numbers: 95.35.+d, 95.55.Ka

I. INTRODUCTION

In light of recent analyses of data from the Fermi Large Area Telescope (LAT) [1] indicating the possibility of an excess of ~ 130 GeV photons from the Galactic Center region [2], there has been renewed interest in dark matter models that can produce monoenergetic photons through the processes $\chi\chi \rightarrow \gamma\gamma, \gamma Z$, and γh [3–9]. Generically, one would expect models with a monoenergetic photon line to be accompanied by a larger continuum photon spectrum. This reasoning derives from general principles: since there are tight constraints on the electromagnetic charge of dark matter, monoenergetic photon production can only arise through a millicharged coupling or through a loop diagram. These processes are expected to be suppressed relative to tree-level annihilation to other Standard Model particles. The dominant annihilation products would then produce a continuum photon flux, either from decays that produce photons or through synchrotron radiation, which is large relative to the strength of a monoenergetic photon signal.

Bounds on the magnitude of the continuum spectrum arising from dark matter annihilation in the Galactic Center thus put tight constraints on dark matter models that could explain a line signal. Many new dark matter models have been proposed to evade these constraints and explain the recent Fermi data [3, 4]. Similarly, a variety of works have studied the question of whether it is possible for the 130 GeV photon excess seen by Fermi to be produced within the MSSM [5–7] or singlet extensions of the MSSM [8], while maintaining consistency with bounds on the continuum spectrum.

In this work, we consider an MSSM analysis from a different perspective. We assume that the MSSM lightest supersymmetric particle (LSP) is the lightest neutralino, χ , and constitutes all of the dark matter in the universe, and we ask what we can learn about the parameters of the MSSM from the observation of a photon line signal and a small (or unobservable) continuum signal. With this goal in mind, we avoid (as far as possible) any assumptions about the MSSM parameter space based on naturalness or simplicity, instead taking as our guide only the mass of the Higgs boson (taken to be $m_h \sim 126$ GeV), other observational constraints, and the assumption of a large photon line signal and small continuum signal. We will find that these demands are so restrictive that they constrain the MSSM parameter space in a way comparable to (though orthogonal to) the typical assumptions that lead to scenarios like the CMSSM or mSUGRA, allowing a scan over allowed parameter space to be feasible.

A large annihilation cross section to monoenergetic photons relative to that for continuum photons can be obtained within the MSSM if the LSP is a purely bino-like neutralino. In this case, the dominant cross section to continuum photons comes from $\chi\chi \rightarrow \tilde{f}f$, a process that is always suppressed in some way. As we demonstrate here, the suppression of the continuum cross section can be large enough that it becomes comparable to the cross section to monoenergetic photons, a scenario at odds with the intuition that tree-level annihilation to continuum photons will have a much larger cross section than loop-suppressed annihilation to final states with monoenergetic photons.

Although this type of analysis is relevant to an understanding of the implications of the Fermi-LAT data, its utility extends beyond that particular observed excess, which may simply be a statistical or systematic effect [10]. Rather, one can consider the implications of the detection of a line signal with future experiments, such as GAMMA-400 [11] or Cherenkov Telescope Array (CTA) [12], or the emergence of such a signal in the data from the Fermi satellite or ground-based gamma-ray detectors such as VERITAS [13] and HESS [14]. The

ground-based CTA will have an effective area several orders of magnitude larger than the effective area of space-based telescopes, but will suffer from the background of cosmic ray-induced Čerenkov showers. The GAMMA-400 satellite, by contrast is not expected to have a significantly larger effective area than the Fermi-LAT, but it may have much better energy resolution. Here we will assume that relevant experiments can distinguish a true line signal from a very hard continuum spectrum, as might arise from a three-body annihilation process such as $\chi\chi \rightarrow f\bar{f}\gamma$.

In section II, we describe the general principles that lead to the region of MSSM parameter space of interest, where the annihilation cross section to monoenergetic photon final states is comparable to that for final states that give rise to a continuum spectrum of photons. In section III we analyze how the cross section for annihilation to various final states scales with relevant model parameters. In section IV we discuss a detailed numerical scan of the annihilation cross sections in this region of parameter space and relevance of our results for more general non-MSSM scenarios. Finally, we state our conclusions in section V.

II. GENERAL PRINCIPLES

As put forward in [7], one can parameterize the limits on dark matter annihilation obtained from gamma-ray observations in terms of two quantities,

$$\begin{aligned}\sigma_{line} &= 2\sigma_{\gamma\gamma} + \sigma_{\gamma Z}, \\ R_{th} &= \frac{\sigma_{ann.}}{\sigma_{line}},\end{aligned}\tag{1}$$

where $\sigma_{ann.}$ is the total dark matter annihilation cross section and $\sigma_{(\gamma\gamma, \gamma Z)}$ are the $\chi\chi \rightarrow (\gamma\gamma, \gamma Z)$ annihilation cross sections, respectively. Since the neutralino LSP is a Majorana fermion, annihilation to the γh final state is p -wave suppressed, and can therefore be ignored. The rate at which dark matter annihilates to monoenergetic photons is thus proportional to σ_{line} , and R_{th} is the ratio of the total annihilation rate to the rate of annihilation to monoenergetic photons. If an excess of monoenergetic photons is observed, then, with some assumptions about the dark matter density profile of the source, one can estimate the quantity $(\sigma_{line}v)$. Assuming the spectra of continuum photons from relevant dark matter annihilation processes are roughly similar in shape, gamma-ray observations can then place an approximate upper bound on R_{th} .

Without any assumptions about the relationships between MSSM parameters, the number of free parameters is so vast that a complete scan is impractical. This consideration usually leads to assumptions motivated by, among other things, naturalness, a string theoretic origin, a putative SUSY breaking mechanism, elegance or simplicity. We will instead consider a framework where the relations assumed between parameters are motivated only by the goal of maximizing $(\sigma_{line}v)$ and minimizing the annihilation to continuum photons. We will find that these considerations are sufficient to drastically reduce the parameter space of the MSSM.

We are guided by general considerations underlying the suppression of annihilation to continuum photons that is often seen in MSSM models.

- Since χ is a Majorana fermion, the cross section for annihilation to Standard Model fermions, $\chi\chi \rightarrow \bar{f}f$ is suppressed either by $v^2 \sim 10^{-6}$ (for p -wave annihilation) or by m_f^2 or the sfermion-mixing angle (for s -wave annihilation). The cross section for annihilation to the light Higgs, $\chi\chi \rightarrow hh$ is also p -wave suppressed.

- In the case of annihilation to fermions, the three-body annihilation processes $\chi\chi \rightarrow \bar{f}f(\gamma, Z, g)$ can be important [15]. These processes are typically suppressed by a factor of α/r^2 or α_s/r^2 , where $r = m_{\tilde{f}}^2/m_f^2$ and $\tilde{f}$ is the exchanged sfermion.
- There is no chirality or p -wave suppression for $\chi\chi \rightarrow W^+W^-, ZZ$, or Zh . But these processes are only allowed at tree-level if χ has some wino or higgsino component; if χ is purely bino, these processes do not occur at tree-level.
- If $m_\chi > m_t$, then the annihilation process $\chi\chi \rightarrow \bar{t}t$ is kinematically allowed. There is no significant suppression of the s -wave annihilation process unless $m_\chi \gg m_t$.

Based on these facts, the low-energy particle content of interest for minimizing R_{th} in the limit where tree-level annihilations to gauge bosons are negligible must include a purely bino-like LSP, negligible sfermion mixing, heavy sbottoms and scharm, and heavy neutral Higgsinos. Although the production of on-shell tops is kinematically forbidden if $m_\chi < m_t$, the production of off-shell tops can still provide a large contribution to (σ_{ann}, v) . Similarly, if $m_\chi \gg m_t$, then although the $\chi\chi \rightarrow \bar{t}t$ annihilation process will be kinematically suppressed, the sensitivity of gamma-ray observations to (σ_{line}, v) will also be suppressed. Finally, we note that if up/down/strange-squarks are light then the $\chi\chi \rightarrow \bar{u}ug, \bar{d}dg$, and $\bar{s}sg$ three-body annihilation processes are only suppressed by α_s . To minimize R_{th} , we will thus assume that all squarks are heavy.

We are therefore led to consider the scenario in which the only sfermions that are possibly light are the sleptons. The charged slepton, sneutrino, and LSP masses are the only parameters that determine both σ_{line} and R_{th} . This scenario is consistent with constraints from the Large Hadron Collider, which has ruled out models with light first generation squarks [17].

III. FEATURES OF THIS SCENARIO

In this section we discuss the general model features that determine the relevant quantities R_{th} and (σ_{line}, v) . Since the LSP is largely bino-like and the squarks and Higgsinos are very heavy, tree-level annihilation occurs largely through t -channel exchange of light sleptons and sneutrinos, as shown in Fig. 1(a). The primary two-body annihilation processes that contribute to (σ_{ann}, v) in this case are $\chi\chi \rightarrow \bar{f}f$ where $f = e, \mu, \tau, \nu_{e,\mu,\tau}$. We parameterize the scale of the various annihilation processes in terms of α , m_χ , and r , as defined above. The cross sections for these processes scale as

$$\propto \left(\frac{\alpha^2}{m_\chi^2 r^2} \right) \times \left[v^2 \text{ or } \left(\frac{m_f^2}{m_\chi^2} \right) \right]. \quad (2)$$

Since $v^2 \approx 10^{-6}$ today, and $m_f^2 \ll m_\chi^2$, these processes are typically negligible.

The relevant three-body annihilation processes are $\chi\chi \rightarrow \bar{l}l(\gamma, Z)$, $\bar{\nu}_l\nu_l Z$, $\bar{l}\nu_l W^-$, and $\bar{\nu}_l l W^+$, where $l = e, \mu, \tau$. These processes occur through t -channel exchange of a light sfermion, with an electroweak gauge boson emitted either from the outgoing fermions or from the internal sfermion line, as shown in Fig. 1(b). If the gauge boson is emitted from the internal line, then the diagram has two scalar propagators and scales as r^{-2} . There are two diagrams in which the gauge boson is emitted from the outgoing fermions, and each diagram has only one scalar propagator. But the diagrams partially cancel, and their sum also scales as r^{-2} [18]. The electroweak bremsstrahlung processes are further suppressed by




FIG. 1. Typical Feynman diagrams for the annihilation processes $\chi\chi \rightarrow \bar{l}l$ (left), $\chi\chi \rightarrow \bar{l}l\gamma$ (center) and $\chi\chi \rightarrow \gamma\gamma$ (right).

an additional factor of α . For $m_\chi \approx \frac{1}{2}m_{Z,W}$, the cross section for annihilation to a three-body state with a massive vector boson will also be phase-space suppressed. Finally, the cross section for annihilation to $\bar{l}l\gamma$ also receives an additional suppression due to angular momentum conservation; for certain regions of the three-body final state phase space, the angular momentum of the final state can only be zero if the gauge boson has helicity zero, which is not possible for a massless gauge boson. If $m_\chi > \frac{1}{2}m_{Z,W}$, then the total three-body annihilation cross section will scale as

$$\propto \left(\frac{\alpha^2}{m_\chi^2 r^2} \right) \times \left[\frac{\alpha}{r^2} \right]. \quad (3)$$

The processes $\chi\chi \rightarrow \gamma\gamma$, γZ proceed through box diagrams with $l, \bar{l}$ running in the loop, as shown in Fig. 1(c). The number of slepton propagators in the loop can vary between one and three, but as the sleptons become heavy, the dominant diagram will have only one slepton in the loop. We thus find

$$(\sigma_{line}v) \propto \left(\frac{\alpha^2}{m_\chi^2 r^2} \right) \times [\alpha^2]. \quad (4)$$

The three-body annihilation process can dominate over $\chi\chi \rightarrow \bar{l}l$ only if $\alpha/r^2 > v^2$. Similarly, three-body annihilation can only dominate over $\chi\chi \rightarrow \bar{b}b$ if $\alpha/r^2 > m_b^2 m_\chi^2 r^2 / m_b^4$. Assuming that the squarks are very heavy, three-body annihilation is the dominant continuum process if $r < \mathcal{O}(10^2)$. As sleptons become heavier, however, annihilations to three-body final states become insignificant, and the two-body final states determine the annihilation cross section to continuum photons, $(\sigma_{cont.}v)$. So we find

$$R_{th} - 1 = \frac{(\sigma_{cont.}v)}{(\sigma_{line}v)} \sim \begin{cases} 1/(\alpha r^2) \sim \mathcal{O}(10^{-2} - 10^2) & \text{for } r \lesssim \mathcal{O}(10^2) \\ v^2/\alpha^2 \sim \mathcal{O}(10^{-2}) & \text{for } r \gtrsim \mathcal{O}(10^2). \end{cases} \quad (5)$$

We thus expect to find values of R_{th} ranging from ~ 1 to ~ 100 . But small R_{th} arises only when the sleptons are heavy, implying that the magnitude of $(\sigma_{line}v)$ will be suppressed. Maximizing the magnitude of the line signal would require $R_{th} \gtrsim \mathcal{O}(100)$.




FIG. 2. Cross section for production of a line signal as a function of dark matter mass, colored by the value of R_{th} : $R_{th} < 5$ (magenta), $5 \leq R_{th} < 10$ (blue), $10 \leq R_{th} < 100$ (cyan), and $R_{th} \geq 100$ (gray).

IV. PARAMETER SPACE SCAN AND ANALYSIS

We calculate the spectrum of supersymmetric particles, Higgs bosons, and dark matter observables using SuSpect [19], FeynHiggs [20], and MicrOMEGAs [21], with supplementary calculations of the annihilation cross sections to three-body final states following [15, 16]. We focus on consistent models that fall at least approximately into the scenario described in section II, and consider the heavy scalars to be degenerate, with masses m_0 in the range $(m_\chi + 3 \text{ TeV}, 8 \text{ TeV})$. We fix the Higgs mixing parameter, μ , to be $\mu = m_0$, to ensure that the lightest neutralino is nearly pure bino, and allow masses for the left- and right-handed¹ $\tilde{e}$, $\tilde{\mu}$, $\tilde{\tau}$, and $\tilde{\nu}_{e,\mu,\tau}$ in the range (m_χ, m_0) . Finally, the trilinear couplings, A , may take values in the range $(-5m_0, 5m_0)$, with larger $|A|$ preferred by the CMS [22] and ATLAS [23] measurements of $m_h \approx 126 \text{ GeV}$. For the parameter points we consider, we require the mass of the lighter CP-even Higgs boson to lie within the range $123 \text{ GeV} < m_h < 129 \text{ GeV}$.

We assume that the lightest neutralino constitutes all of the dark matter in the universe. In figure 2, we plot the cross section for production of a line signal, $(\sigma_{line}v)$, as a function of dark matter mass. The points are color-coded by the value of R_{th} : $R_{th} < 5$ (magenta), $5 \leq R_{th} < 10$ (blue), $10 \leq R_{th} < 100$ (cyan), and $R_{th} \geq 100$ (gray). In the left panel, we address potential line signals observable by the Fermi satellite, for neutralino masses up to roughly 300 GeV, while in the right panel we consider neutralino masses as large as 2.5 TeV. Though HESS and VERITAS are sensitive to dark matter masses in this range, the cross sections to monoenergetic photons are all well below the current limits [14]. We note that, with adequate energy resolution, the $\gamma\gamma$ and γZ line signals will be distinguishable, and R_{th} and $(\sigma_{line}v)$ will not be optimal parametrizations of the signal strength. However any observed line signal(s) can be simply mapped to particular values of R_{th} and $(\sigma_{line}v)$.

Finally, the largest cross sections, $(\sigma_{line}v) \approx 10^{-29} \text{ cm}^3/\text{s}$, are indeed obtained in the limit $m_{\tilde{l}} \approx m_\chi$, the limit discussed in Ref. [3]. GAMMA-400 may be sensitive to cross sections of this size for small enough m_χ [24]. It has been shown that for $m_{\tilde{l}} \approx m_\chi$, very hard bremsstrahlung photons can mimic a gamma-ray line signal for the present Fermi-LAT energy resolution [5]. Here, all electroweak bremsstrahlung cross sections are included as

¹ By “left- and right-handed” scalars, we mean the partners of the left- and right-handed Standard Model leptons.




FIG. 3. The cross section for production of a line signal as a function of dark matter mass, colored by the value of R_{th} : $R_{th} < 5$ (magenta), $5 \leq R_{th} < 10$ (blue), $10 \leq R_{th} < 100$ (cyan), and $R_{th} \geq 100$ (gray). Here, the thermal relic abundance of neutralinos is restricted to be $\Omega_\chi h^2 < 0.15$.

part of the annihilation cross section to continuum photons. Therefore one can interpret R_{th} for the points with the largest cross sections in figure 2 as maximal, since any very hard bremsstrahlung will effectively contribute to a line signal, as perceived by Fermi-LAT, and not to the perceived continuum.

From the analysis of the previous section, we expect that models with small $(\sigma_{line}v)$ have large values of r , which in turn implies small R_{th} . This expectation is borne out in figure 2 for $(\sigma_{line}v) \gtrsim 10^{-31} \text{ cm}^3/\text{s}$. However, as r becomes very large, the decoupling approximation no longer applies to the parameter scan (since the masses of the sleptons, squarks, neutralinos and charginos become comparable), and annihilations to final states that produce continuum photons become much more significant than annihilations to final states with monoenergetic photons. We note that the dearth of models with small R_{th} and very small $(\sigma_{line}v)$ (see the left panel of figure 2) is an artifact of the imposed cut-off $m_{\tilde{l}} \leq m_0 \leq 8 \text{ TeV}$.

In figure 3 we focus on neutralinos with $m_\chi \leq 300 \text{ GeV}$ and plot only points for which the thermal relic abundance of dark matter satisfies $\Omega_\chi h^2 < 0.15$, in rough compatibility with the observed abundance of cold dark matter [25]. We see that some of the very largest line cross sections are due to models in which the dark matter is a thermal relic. It is not surprising that we find points for which $R_{th} \sim \mathcal{O}(10^2)$ with $(\sigma_{line}v) \sim 10^{-29} \text{ cm}^3/\text{s}$, i.e. a continuum cross section of $(\sigma_{cont.v}) \sim 10^{-27} \text{ cm}^3/\text{s}$, allowing compatibility between the measured dark matter abundance and the thermal relic density. We also find points with both smaller $(\sigma_{line}v)$ and smaller R_{th} . Indeed, there are even points for which $R_{th} < 5$ that can predict a relic abundance of neutralinos compatible with the observed dark matter abundance. Unfortunately, most of these points have $(\sigma_{line}v) \sim 10^{-29} - 10^{-30} \text{ cm}^3/\text{s}$, below the reach of the Fermi-LAT, and likely below the reach of GAMMA-400, as well [24]. For these models, the annihilation cross section at dark matter freeze-out was dominated by terms that are suppressed in the present era (for example, p -wave terms).

Although this analysis has been presented in the framework of the MSSM, it applies more generally to any model where the dark matter candidate is a Majorana fermion that is neutral under the Standard Model gauge groups. For such a model, dark matter can annihilate at tree-level only to $\bar{f}f$ or hh , with the latter process being either p -wave suppressed or suppressed by CP-violating phases. If annihilation to $\bar{f}f$ proceeds through t -channel exchange of a scalar partner, then the analysis given here for the relation between R_{th} and the mass

of the scalar partner still holds. Note, however, that the bino-fermion-sfermion coupling is simply related to the hypercharge coupling. For a more general dark matter model, this coupling, g , is a free parameter. Since $(\sigma_{line}v)$ and all continuum annihilation cross sections scale as g^4 , the choice of coupling does not affect R_{th} . However, it does affect the overall scale of the annihilation cross section; if this coupling is required to be perturbative, then $\sigma_{(\gamma\gamma,\gamma Z)}$ can be increased by up to 4 orders of magnitude, in agreement with the limit discussed in Ref. [3]. For models with small R_{th} , such an enhancement could bring a line signal within the reach of GAMMA-400.

V. CONCLUSIONS

We have considered a class of MSSM models for which the ratio $\sigma(\chi\chi \rightarrow \gamma\gamma, \gamma Z)/\sigma(\chi\chi \rightarrow \text{anything})$ is maximized. We have found that this requirement leads to a dramatic restriction in the number of relevant MSSM parameters. In particular, the region of interest studied here contains a nearly pure bino LSP, with the only light superpartners being the bino and the sleptons. Interestingly, this region of parameter space is consistent with recent bounds from the LHC, which tightly constrain models with light squarks.

For such models, we find that if a photon line signal is potentially observable at current or next generation experiments, then R_{th} can be as low as $\mathcal{O}(10 - 100)$. Moreover, for many such models the continuum signal is dominated by three-body annihilation. Since three-body annihilation may produce a very hard photon spectrum, it is possible that such a continuum annihilation process could be mistaken for a line. Therefore our numerical results for R_{th} represent the maximal values possible (i.e. the limit of perfect energy resolution). The improved energy resolution of next generation gamma-ray telescopes should make it possible to resolve this difference.

For MSSM models with small R_{th} , the line signal will be a few orders of magnitude too small to be observed with Fermi. However, more general models of Standard Model-neutral Majorana fermion dark matter can achieve the same R_{th} with an enhancement in the line signal of up to four orders of magnitude, potentially bringing a pure line signal with no observable continuum spectrum within reach of observation.

Acknowledgments

We thank the organizers of ICHEP2012 and the Center for Theoretical Underground Physics and Related Areas (CETUP* 2012) in South Dakota for their support and hospitality during the completion of this work. We are grateful to D. Marfatia and C. Kelso for useful discussions, and to K. Fukushima for comments and for assistance with the numerical code for computing the continuum annihilation cross section. Support and resources from the Center for High Performance Computing at the University of Utah are also gratefully acknowledged. The work of J. K. is supported in part by Department of Energy grant DE-FG02-04ER41291.

[1] <http://www-glast.stanford.edu>

[2] T. Bringmann, X. Huang, A. Ibarra, S. Vogl and C. Weniger, JCAP **1207**, 054 (2012); [arXiv:1203.1312 [hep-ph]]; C. Weniger, JCAP **1208**, 007 (2012) [arXiv:1204.2797 [hep-ph]];

- E. Tempel, A. Hektor and M. Raidal, JCAP **1209**, 032 (2012) [Addendum-ibid. **1211**, A01 (2012)]; M. Su and D. P. Finkbeiner, arXiv:1206.1616 [astro-ph.HE].
- [3] M. R. Buckley and D. Hooper, Phys. Rev. D **86**, 043524 (2012) [arXiv:1205.6811 [hep-ph]];
- [4] E. Dudas, Y. Mambrini, S. Pokorski and A. Romagnoni, JHEP **1210** (2012) 123 [arXiv:1205.1520 [hep-ph]]; J. M. Cline, Phys. Rev. D **86** (2012) 015016 [arXiv:1205.2688 [hep-ph]]; W. Buchmuller and M. Garny, JCAP **1208**, 035 (2012) [arXiv:1206.7056 [hep-ph]]; M. T. Frandsen, U. Haisch, F. Kahlhoefer, P. Mertsch and K. Schmidt-Hoberg, JCAP **1210**, 033 (2012) [arXiv:1207.3971 [hep-ph]]; J. -C. Park and S. C. Park, Phys. Lett. B **718**, 1401 (2013) [arXiv:1207.4981 [hep-ph]]; S. Tulin, H. -B. Yu and K. M. Zurek, arXiv:1208.0009 [hep-ph]; L. Bergstrom, Phys. Rev. D **86**, 103514 (2012) [arXiv:1208.6082 [hep-ph]]; L. Wang and X. -F. Han, Phys. Rev. D **87**, 015015 (2013) [arXiv:1209.0376 [hep-ph]]; J. Fan and M. Reece, arXiv:1209.1097 [hep-ph]; N. Weiner and I. Yavin, Phys. Rev. D **87**, 023523 (2013) [arXiv:1209.1093 [hep-ph]]; H. M. Lee, M. Park and W. -I. Park, JHEP **1212**, 037 (2012) [arXiv:1209.1955 [hep-ph]]; S. Baek, P. Ko and E. Senaha, arXiv:1209.1685 [hep-ph]; M. R. Buckley and E. T. Neil, Phys. Rev. D **87**, 043510 (2013) [arXiv:1209.6054 [hep-ph]]; F. D'Eramo, M. McCullough and J. Thaler, arXiv:1210.7817 [hep-ph]; K. Schmidt-Hoberg, F. Staub and M. W. Winkler, [arXiv:1211.2835 [hep-ph]]; Y. Farzan and A. R. Akbarieh, arXiv:1211.4685 [hep-ph]; Y. Bai, M. Su and Y. Zhao, [arXiv:1212.0864 [hep-ph]]; Y. Zhang, arXiv:1212.2730 [hep-ph]; A. Biswas, D. Majumdar, A. Sil and P. Bhattacharjee, arXiv:1301.3668 [hep-ph]; P. -H. Gu, arXiv:1301.4368 [hep-ph]. J. Chen and Y. -F. Zhou, arXiv:1301.5778 [hep-ph]; C. B. Jackson, G. Servant, G. Shaughnessy, T. M. P. Tait and M. Taoso, arXiv:1302.1802 [hep-ph]; A. Rajaraman, T. M. P. Tait and A. M. Wijangco, arXiv:1211.7061 [hep-ph]; A. Rajaraman, T. M. P. Tait and D. Whiteson, JCAP **1209**, 003 (2012) [arXiv:1205.4723 [hep-ph]].
- [5] B. Shakya, arXiv:1209.2427 [hep-ph].
- [6] W. Buchmuller and M. Garny, JCAP **1208**, 035 (2012) [arXiv:1206.7056 [hep-ph]]; T. Bringmann and C. Weniger, Phys. Dark Univ. **1**, 194 (2012) [arXiv:1208.5481 [hep-ph]].
- [7] T. Cohen, M. Lisanti, T. R. Slatyer and J. G. Wacker, arXiv:1207.0800 [hep-ph].
- [8] G. Chalons, M. J. Dolan and C. McCabe, JCAP **1302**, 016 (2013) [arXiv:1211.5154 [hep-ph]]; D. Das, U. Ellwanger and P. Mitropoulos, JCAP **1208**, 003 (2012) [arXiv:1206.2639 [hep-ph]]; Z. Kang, T. Li, J. Li and Y. Liu, arXiv:1206.2863 [hep-ph]; J. Kozaczuk, S. Profumo and C. L. Wainwright, arXiv:1302.4781 [hep-ph].
- [9] E. Dudas, Y. Mambrini, S. Pokorski and A. Romagnoni, JHEP **0908** (2009) 014 [arXiv:0904.1745 [hep-ph]]; Y. Mambrini, JCAP **0912** (2009) 005 [arXiv:0907.2918 [hep-ph]].
- [10] D. Whiteson, JCAP **1211**, 008 (2012) [arXiv:1208.3677 [astro-ph.HE]]; A. Hektor, M. Raidal and E. Tempel, arXiv:1209.4548 [astro-ph.HE]; D. P. Finkbeiner, M. Su and C. Weniger, JCAP **1301**, 029 (2013) [arXiv:1209.4562 [astro-ph.HE]]; D. Whiteson, arXiv:1302.0427 [astro-ph.HE].
- [11] A. M. Galper, O. Adriani, R. L. Aptekar, I. V. Arkhangelskaja, A. I. Arkhangel'skiy, M. Boezio, V. Bonvicini and K. A. Boyarchuk *et al.*, Adv. Space Res. **51**, 297 (2013) [arXiv:1201.2490 [astro-ph.IM]]; A. M. Galper, O. Adriani, R. L. Aptekar, I. V. Arkhangelskaja, A. I. Arkhangel'skiy, M. Boezio, V. Bonvicini and K. A. Boyarchuk *et al.*, AIP Conf. Proc. **1516**, 288 (2012) [arXiv:1210.1457 [astro-ph.IM]].
- [12] M. Actis *et al.* [CTA Consortium Collaboration], Exper. Astron. **32**, 193 (2011) [arXiv:1008.3703 [astro-ph.IM]].
- [13] author = R. Ong *et al.*, booktitle = in proceedings of 31st ICRC, Lodz, Poland, astro-

- ph/912.5355, year = 2009.
- [14] A. Abramowski *et al.* [H.E.S.S. Collaboration], Phys. Rev. Lett. **110**, 041301 (2013) [arXiv:1301.1173 [astro-ph.HE]]; Y. Becherini *et al.* [HESS Collaboration], AIP Conf. Proc. **1505**, 741 (2012).
 - [15] V. Barger, W. -Y. Keung and D. Marfatia, Phys. Lett. B **707**, 385 (2012) [arXiv:1111.4523 [hep-ph]]; N. F. Bell, J. B. Dent, A. J. Galea, T. D. Jacques, L. M. Krauss and T. J. Weiler, Phys. Lett. B **706**, 6 (2011) [arXiv:1104.3823 [hep-ph]]; L. Bergstrom, T. Bringmann and J. Edsjo, Phys. Rev. D **78**, 103520 (2008) [arXiv:0808.3725 [astro-ph]]; M. Garny, A. Ibarra and S. Vogl, JCAP **1107**, 028 (2011) [arXiv:1105.5367 [hep-ph]]; M. Garny, A. Ibarra and S. Vogl, arXiv:1112.5155 [hep-ph]; N. F. Bell, A. J. Brennan and T. D. Jacques, “Neutrino signals from electroweak bremsstrahlung in solar WIMP annihilation,” arXiv:1206.2977 [hep-ph].
 - [16] K. Fukushima, Y. Gao, J. Kumar and D. Marfatia, Phys. Rev. D **86**, 076014 (2012) [arXiv:1208.1010 [hep-ph]].
 - [17] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Rev. Lett. **107**, 221804 (2011) [arXiv:1109.2352 [hep-ex]]; G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **710**, 67 (2012) [arXiv:1109.6572 [hep-ex]].
 - [18] P. Ciafaloni, M. Cirelli, D. Comelli, A. De Simone, A. Riotto and A. Urbano, JCAP **1106**, 018 (2011) [arXiv:1104.2996 [hep-ph]].
 - [19] A. Djouadi, J. -L. Kneur and G. Moultaka, Comput. Phys. Commun. **176**, 426 (2007) [hep-ph/0211331].
 - [20] T. Hahn, S. Heinemeyer, W. Hollik, H. Rzehak and G. Weiglein, Nucl. Phys. Proc. Suppl. **205-206**, 152 (2010) [arXiv:1007.0956 [hep-ph]].
 - [21] G. Belanger, F. Boudjema, P. Brun, A. Pukhov, S. Rosier-Lees, P. Salati and A. Semenov, Comput. Phys. Commun. **182**, 842 (2011) [arXiv:1004.1092 [hep-ph]].
 - [22] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012).
 - [23] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012).
 - [24] L. Bergstrom, G. Bertone, J. Conrad, C. Farnier and C. Weniger, JCAP **1211**, 025 (2012) [arXiv:1207.6773 [hep-ph]].
 - [25] G. Hinshaw, D. Larson, E. Komatsu, D. N. Spergel, C. L. Bennett, J. Dunkley, M. R. Nolta and M. Halpern *et al.*, arXiv:1212.5226 [astro-ph.CO].